

**A. Brauer, P. Dulski, C. Mangili, J. Mingram and J. Liu**

- Alley, R.B. et al., 2003: Abrupt climate change, *Science*, **299**: 2005-2010.
- Brauer, A., 2004: Annually laminated lake sediments and their palaeoclimatic relevance. In: Fischer, H., et al. (Eds), *The Climate in Historical Times. Towards a Synthesis of Holocene Proxy Data and Climate Models*, Springer Verlag, 109-127.
- Brauer, A., Allen, J.R.M., Mingram, J., Dulski, P., Wulf, S. and Huntley, B., 2007: Evidence for last interglacial chronology and environmental change from Southern Europe, *Proceedings of the National Academy of Science*, **104**: 450-455.
- Brauer, A., Haug, G.H., Dulski, P., Sigman, D.M. and Negendank, J.F.W., 2008a: An abrupt wind shift in Western Europe at the onset of the Younger Dryas cold period, *Nature Geoscience*, **1**: 520-523.
- Brauer, A., Mangili, C., Moscariello, A. and Witt, A., 2008b: Palaeoclimatic implications from micro-facies data of a 5900 varve time series from the Pianico interglacial sediment record, Southern Alps, *Palaeogeography Palaeoclimatology, Palaeoecology*, **259**: 121-135.
- Broecker, W.S., 2006: Was the Younger Dryas triggered by a flood? *Science*, **312**: 1146 - 1148.
- Chu, G., Liu, J., Schettler, G., Li, J., Sun, Q., Gu, Z., Lu, H., Liu, Q. and Liu, T., 2005: Sediment Fluxes and Varve Formation in Sihailongwan, a Maar Lake from Northeastern China, *Journal of Paleolimnology*, **34**: 311-324.
- Kitagawa, H. and van der Plicht, J., 1998: Atmospheric radiocarbon calibration to 45,000 yr B.P.: Late Glacial fluctuations and cosmogenic isotope production, *Science*, **279**: 1187-1190.
- Lamoureux, S.F., 2001: Varve chronology techniques. In: Last, W.M. and Smol, J.P. (Eds), *Developments in Paleoenvironmental Research (DPER), Vol. 2 - Tracking Environmental Change Using Lake Sediments: Physical and Chemical Techniques*, Kluwer, Dordrecht, 247-260.
- Lotter, A.F. and Lemcke, G., 1999: Methods for preparing and counting biochemical varves, *Boreas*, **28**: 243-252.
- Mangili, C., Brauer, A., Moscariello, A. and Naumann, R., 2005: Microfacies of detrital event layers deposited in Quaternary varved lake sediments of the Piànico-Sèllere Basin (northern Italy). *Sedimentology*, **52**, 927-943.
- Mangili, C., Brauer, A., Plessen, B. and Moscariello, A., 2007: Centennial-scale oscillations in stable oxygen and carbon isotopes of endogenic calcite from a 15500 varve year record of the Piànico interglacial, *Quaternary Science Reviews*, **26**: 1725-1735.
- Merkt, J., 1971: Zuverlässige Auszählungen von Jahresschichten in Seesedimenten mit Hilfe von Groß-Dünnschliffen, *Archiv für Hydrobiologie*, **69**: 145-154.
- Mingram, J., Allen, J.R.M., Brüchmann, C., Liu, J., Luo, X., Negendank, J.F.W., Nowaczyk N. and Schettler, G., 2004: Maar- and crater lakes of the Long Gang Volcanic Field (N.E. China) - overview, laminated sediments, and vegetation history of the last 900 years, *Quaternary International*, **123-125**: 135-147.
- Rioual, P., Chu, G., Han, J., Mingram, J., Liu, Q., Schettler, G. and Liu, J., 2006: Holocene climatic and limnologic changes at Sihailongwan maar lake (Northeastern China) as indicated by diatom remains: preliminary results, *HOLIVAR Natural Climate Variability and Global Warming*, London.
- Schettler, G., Mingram, J., Liu, Q., Stebich M. and Dulski, P., 2006: East-Asian monsoon variability between 15000 and 2000 cal. yr BP recorded in varved sediments of Lake Sihailongwan (northeastern China, Long Gang volcanic field), *The Holocene*, **16**: 1043-1057.
- Steffensen, J.P. et al., 2008: High-Resolution Greenland Ice Core Data Show Abrupt Climate Change Happens in Few Years, *Science*, **321**: 680-684.

- Zolitschka, B. 2003: Dating based on freshwater and marine laminated sediments. In: MacKay, A. et al. (Eds), *Global Change in the Holocene*, Edward Arnold Publishers, London: 92-106.
- Zolitschka, B. 2006: Varved lake sediments. In: Elias, S.A. (Ed.), *Encyclopedia of Quaternary Science*, Amsterdam, Elsevier, 3105-3114.

**M.J.L. Coolen and J.A.E. Gibson**

- Bissett, A., Gibson, J.A.E., Jarman, S.N., Swadling, K.M. and Cromer, L., 2005: Isolation, amplification, and identification of ancient copepod DNA from lake sediments, *Limnology and Oceanography: Methods*, **3**: 533-542.
- Boere, A., Abbas, B., Rijpstra, W.I.C., Versteegh, G.J.M., Volkman, J.K., Sinninghe Damsté, J.S. and Coolen, M.J.L., 2009: Late-Holocene succession of dinoflagellates in an Antarctic fjord using a multi-proxy approach: paleoenvironmental genomics, lipid biomarkers, and palynomorphs, *Geobiology* **7**: 265-281.
- Brassell, S.C., Eglinton, G., Marlowe, I.T., Pflaumann, U. and Sarnthein, M., 1986: Molecular stratigraphy: A new tool for climatic assessment, *Nature*, **320**: 129-133.
- Coolen, M.J.L. and Overmann, J., 1998: Analysis of subfossil molecular remains of purple sulfur bacteria in a lake sediment, *Applied and Environmental Microbiology*, **64**: 4513-4521.
- Coolen, M.J.L. and Overmann, J., 2007: 217 000-year-old DNA sequences of green sulfur bacteria in Mediterranean sapropels and their implications for the reconstruction of the paleoenvironment, *Environmental Microbiology*, **9**: 238-249.
- Coolen, M.J.L. and Shtereva, G., 2009: Vertical distribution of metabolically active eukaryotes in the water column and sediments of the Black Sea, *FEMS Microbial Ecology* (Accepted for publication).
- Coolen, M.J.L., Muyzer, G., Rijpstra, W.I.C., Schouten, S., Volkman, J.K. and Sinninghe Damsté, J.S., 2004: Combined DNA and lipid analyses of sediments reveal changes in Holocene haptophyte and diatom populations in an Antarctic lake, *Earth and Planetary Science Letters*, **223**: 225-239.
- Coolen, M.J.L., Muyzer, G., Schouten, S., Volkman, J.K. and Sinninghe Damsté, J.S., 2006: Sulfur and methane cycling during the Holocene in Ace Lake (Antarctica) revealed by lipid and DNA stratigraphy. In: Neretin, L.N. (ed), *Past and Present Marine Water Column Anoxia*, NATO Science Series: IV-Earth and Environmental Sciences, Springer, 41-65.
- Coolen, M.J.L., Saenz, J.P., Giosan, L., Trowbridge, N.Y., Dimitrov, P., Dimitrov, D., and Eglinton, T.I., 2009: DNA and lipid molecular stratigraphic records of haptophyte succession in the Black Sea during the Holocene. *Earth Planet Sc Lett* **284**: 610-621.
- Coolen, M.J.L., Volkman, J.K., Abbas, B., Muyzer, G., Schouten, S. and Sinninghe Damsté, J.S., 2007: Identification of organic matter sources in sulfidic late Holocene Antarctic fjord sediments from fossil rDNA sequence analysis, *Paleoceanography*, **22**: PA2211, doi:10.1029/2006PA001309.
- Coolen, M.J.L., Talbot, H.M., Abbas, B.A., Ward, C., Schouten, S., Volkman, J.K. and Sinninghe Damsté, J.S., 2008: Sources for sedimentary bacteriohopanepolyols as revealed by 16S rDNA stratigraphy, *Environmental Microbiology*, **10**: 1783-1803.
- Cromer, L., Gibson, J. A. E., Swadling, K. M. and Ritz, D.A., 2005: Faunal microfossils: Indicators of Holocene ecological change in a saline Antarctic lake, *Palaeogeography, Palaeoclimatology, Palaeoecology*, **221**: 83-97.
- D'Andrea, W.J., Lage, M., Martiny, J.B.H., Laatsch, A.D., Amaral-Zettler, L.A., Sogin, M.L. and Huang, Y., 2006: Alkenone producers inferred from well-preserved 18S rDNA in Greenland lake sediments, *Journal of Geophysical Research*, **111**: G0313: doi:10.1029/2005JG000121.

- Epp, L., Stoof, K., Trauth, M.H. and Tiedemann, R., 2009: Historical genetics on a sediment core from a Kenyan lake: Intraspecific genotype turnover in a tropical rotifer is related to past environmental changes, accepted in *Journal of Paleolimnology*.
- Gibson, J.A.E. and Bayly, I.A.E., 2007: New insights into the origins of crustaceans of Antarctic lakes, *Antarctic Science*, **19**: 157-164.
- Manske, A.K., Henssge, U., Glaeser, J. and Overmann, J., 2008: Subfossil 16S rRNA gene sequences of green sulfur bacteria in the Black Sea and their implications for past photic zone anoxia, *Applied and Environmental Microbiology*, **74**: 624-632.
- Willerslev, E. et al., 2007: Ancient biomolecules from deep ice cores reveal a forested Southern Greenland, *Science*, **317**: 111-114.

**P. Francus, H. Lamb, T. Nakagawa, M. Marshall, E. Brown and Suigetsu 2006 Project members**

- Brown, E.T., Johnson, T.C., Scholz, C.A., Cohen, A.S. and King, J.W., 2007: Abrupt Change in Tropical African Climate Linked to the Bipolar Seesaw Over the Past 55,000 Years, *Geophysical Research Letters*, **34**: doi:10.1029/2007GL031240.
- Colman, S.M., Peck, J.A., Karabanov, E.B., Carter, S.J., Bradbury, J.P., King, J.W. and Williams, D. F., 1995: Continental climate response to orbital forcing from biogenic silica records in Lake Baikal, *Nature*, **378**: 769-771.
- Croudace, I.W., Rindby, A. and Rothwell, R.G., 2006: ITRAX: description and evaluation of a new X-ray core scanner. In: Rothwell, R.G., (Ed.) *New ways of looking at sediment cores and core data*, Geological Society Special Publication, **267**: 51-63.
- DeMaster, D.J., 1979: *The Marine Budgets of Silica and <sup>32</sup>Si*, PhD thesis, Yale University.
- Guyard, H., Chapron, E., St-Onge, G., Anselmetti, F., Arnaud, F., Magand, O., Francus, P. and Mélières, M.-A., 2007: High-altitude varve records of abrupt environmental changes and mining activity since the Bronze Age in the Western French Alps (Lake Bramant, Grandes Rousses Massif), *Quaternary Science Reviews*, **26**: 2644–2660.
- Jansen, J.H.F., Van der Gaast, S.J., Koster, B. and Vaars, A.J., 1998: CORTEX, a shipboard XRF-scanner for element analyses in split sediment cores, *Marine Geology*, **151**: 143-153.
- Johnson, T.C., Brown, E.T., McManus, J., Barry, S., Barker, P. and Gasse, F., 2002: A high-resolution paleoclimate record spanning the past 25,000 years in Southern East Africa, *Science*, **296**: 113-132.
- Nakagawa, T., 2007: Double-L channel: an amazingly non-destructive method of continuous sub-sampling from sediment cores, *Quaternary International*, **167-168**: Supplement: 298.
- Rasmussen, S.O. et al., 2006: A new Greenland ice core chronology for the last glacial termination, *Journal of Geophysical Research*, **111**: D06102, doi:10.1029/2005JD006079.
- Rosman, K.J.R., Chisholm, W., Hong, S., Candelone, J.P. and Boutron, C.F., 1997: Lead from Carthaginian and Roman Spanish mines isotopically identified in Greenland ice dated from 600 BC to 300 AD, *Environmental Science & Technology*, **31**: 3413–3416.
- St-Onge, G., Mulder, T., Francus, P. and Long, B., 2007: Continuous physical properties of cored marine sediments. In Hillaire-Marcel, C. and de Vernal, A., (Eds), *Proxies in late Cenozoic Paleoceanography*, Elsevier, 63-98.
- Weltje, G.J. and Tjallingii, R., 2008. Calibration of XRF core scanners for quantitative geochemical logging of sediment cores: Theory and application, *Earth and Planetary Science Letters*, **274**: 423-438.

**F. García-Rodríguez, E. Piovano, L. del Puerto, H. Inda, S. Stutz, R. Bracco, D. Panario, F. Córdoba, F. Sylvestre and D. Ariztegui**

- Angulo, R.J, Lessa, G.C. and de Souza, M.C., 2006: A critical review of mid-to late Holocene sea-level fluctuations on the eastern Brazilian coastline, *Quaternary Science Reviews*, **25**: 486-506.
- Bracco, R., Inda, H., del Puerto, L., Castiñeira, C, Sprechmann, P. and García-Rodríguez, F., 2005: Relationships between Holocene sea level variation, trophic development and climate change in Negra Lagoon, southern Uruguay, *Journal of Paleolimnology*, **33**: 252-262.
- Bracco, R., García-Rodríguez, F., del Puerto, L., Inda, H. and Castiñeira, C., 2008: Holocene relative sea-level variation inferred from records in the basin of Castillos. In: *Structure and function of world shallow lakes*, Book of Abstracts, Punta del Este, Uruguay. p. 75.
- Cavallotto, J.L., Violante, R.A. and Parker, G., 2004: Sea-level fluctuations during the last 8600 years in the de la Plata River (Argentina), *Quaternary International*, **114**: 155–165.
- del Puerto L., 2009: *Silicofitolitos como Indicadores Paleoambientales: bases comparativas y reconstrucción paleoclimática a partir del Pleistoceno Tardío en el SE del Uruguay*, M.Sc. Thesis PEDECIBA, Facultad de Ciencias, Universidad de la República, Montevideo, 189.
- del Puerto, L., Inda, H. and García-Rodríguez, F., 2008: Reconstrucción Paleoambiental para el Holoceno medio y tardío en la cuenca de la Laguna Negra: el aporte de los indicadores biosilíceos. In: Korstanje, A. and Babot, P. (Eds), *Matices Interdisciplinarios en Estudios Fitolíticos y de Otros Microfósiles*, BAR International Series 1870: 119-129.
- del Puerto, L, García-Rodríguez, F., Inda H., Bracco, R., Castiñeira, C., J.B. and Adams, J.B., 2006: Paleolimnological evidence of Holocene paleoclimatic changes in Lake Blanca, southern Uruguay, *Journal of Paleolimnology*, **36**: 151-163.
- García-Rodríguez, F., Mazzeo N, Sprechmann P., Metzeltin D., Sosa F., Treutler H.C., Renom M., Scharf B. and Gaucher C., 2002: Paleolimnological assessment of human impacts in Lake Blanca, SE Uruguay, *Journal of Paleolimnology*, **28**: 457-468.
- García-Rodríguez, F. et al., 2004: Holocene trophic state changes in relation to sea level variation in Lake Blanca, SE Uruguay, *Journal of Paleolimnology*, **31**: 99-115.
- Inda, H., 2009: Paleolimnología de cuerpos de agua someros del sudeste del Uruguay: evolución holocénica e impacto humano, M.Sc. Thesis. PEDECIBA-Biología, Universidad de la República.
- Inda, H. et al., 2008: Mid-Holocene palaeoenvironmental reconstruction and recent human impact in Laguna del Diario, SE Uruguay. In: *Structure and function of world shallow lakes*, Book of Abstracts, Punta del Este, Uruguay, 77.
- Isla, F., 1998: Holocene coastal evolution of Buenos Aires, *Quaternary of South America and Antarctic Peninsula*, **11**: 297-321
- Martin, L. and Suguio, K., 1992: Variation of coastal dynamics during the last 7000 years recorded in beachridge plains associated with river mouths: example from the Central Brazilian Coast, *Palaeogeography, Palaeoclimatology, Palaeoecology*, **99**: 119-140.
- Metzeltin, D., Lange-Bertalot, H. and García-Rodríguez, F., 2005: Diatoms of Uruguay. In: H. Lange-Bertalot (Ed.). *Iconographia Diatomologica*, Vol. 15. A.R.G. Gantner Verlag, Koeltz Scientific Books, Königstein, 737 pp.
- Metzeltin, D. and García Rodríguez, F., 2003: *Las Diatomeas Uruguayas*, DIRAC Ediciones, Facultad de Ciencias, Montevideo, Uruguay, 208 pp.
- Piovano, E., Ariztegui, D. and Damatto Moreira, S., 2002: Recent environmental changes in Laguna Mar Chiquita (Central Argentina): A sedimentary model for a highly variable saline lake, *Sedimentology*, **49**: 1371-1384.
- Smol, J.P., 1985: The ratio of diatom frustules to chrysophycean statospores: a useful paleolimnological index, *Hydrobiologia*, **123**: 199-208.
- Vera, C. et al., 2006: Towards a unified view of the American Monsoon System, *Journal of Climate*, **19**: 4977-5000.

**M. Grosjean, L. von Gunten, M. Trachsel and C. Kamenik**

- Appleby, P.G., 2008: Three decades of dating recent sediments by fallout radionuclides: a review, *Holocene*, **18**: 83–93.
- Arnaud, F., Lignier, V., Revel, M., Desmet, M., Beck, C., Pourchet, M., Charlet, F., Trentesaux, A. and Tribovillard, N., 2002: Flood and earthquake disturbance of Pb-210 geochronology (Lake Anterne, NW Alps), *Terra Nova*, **14**: 225–232.
- Birks, H.J.B., 1998: Numerical tools in paleolimnology – Progress, potentials, and problems, *Journal of Paleolimnology*, **20**: 307–332.
- Carroll, J. and Lerche, I., 2003: *Sedimentary processes: quantification using radionuclides*, Elsevier, Oxford.
- Cook, E.R., Briffa, K.R. and Jones, P.D., 1994: Spatial regression methods in dendroclimatology: A review and comparison of two techniques, *International Journal of Climatology*, **14**: 379–402.
- Esper, J., Frank, D.C., Wilson, R.J.S. and Briffa, K.R., 2005: Effect of scaling and regression on reconstructed temperature amplitude for the past millennium, *Geophysical Research Letters*, **32**.
- Kalugin, I., Daryin, A., Smolyaninova, L., Andreev, A., Diekmann, B. and Khlystov, O., 2007: 800-yr-long records of annual air temperature and precipitation over southern Siberia inferred from Teletskoye Lake sediments, *Quaternary Research*, **67/3**: 400–410.
- Koinig, K.A. and 11 co-authors, 2002: Environmental changes in an alpine lake (Gossenköllesee, Austria) over the last two centuries—the influence of air temperature on biological parameters, *Journal of Paleolimnology*, **28**: 147–160.
- Rein, B. and Sirocko, F., 2002: In-situ reflectance spectroscopy - analysing techniques for high-resolution pigment logging in sediment cores, *International Journal of Earth Sciences*, **91**: 950–954.
- Trachsel, M., Eggenberger, U., Grosjean, M., Blass, A. and Sturm, M., 2008: Mineralogy-based quantitative precipitation and temperature reconstructions from annually laminated lake sediments (Swiss Alps) since AD 1580, *Geophysical Research Letters*, **35(13)**: L13707.
- von Gunten, L., Grosjean, M., Beer, J., Grob, P., Morales, A. and Urrutia, R., 2009a: Age modeling of young non-varved lake sediments: methods and limits. Examples from two lakes in Central Chile, *Journal of Paleolimnology*, DOI 10.1007/s10933-008-9284-5.
- von Gunten, L., Grosjean, M., Rein, B., Urrutia, R. and Appleby, P., 2009: A quantitative high-resolution summer temperature reconstruction based on sedimentary pigments from Laguna Aculeo, central Chile, back to AD 850, *The Holocene*, **19/6**: 873–881.

**P. Gell, H. Bennion and R. Battarbee**

- Battarbee, R.W. 1990: The causes of lake acidification, with special reference to the role of acid deposition, *Philosophical Transactions of the Royal Society, London B*, **327**: 339–347.
- Battarbee, R.W., Morley, D., Bennion, H. and Simpson, G.L., 2007: A meta-database for recent paleolimnological studies, *PAGES News*, **15**: 23–24.
- Bennion, H. and Battarbee R., 2007: The European Union Water Framework Directive: opportunities for palaeolimnology, *Journal of Paleolimnology*, **38**: 285–295.
- Bennion, H., Fluin, J. and Simpson, G.L. 2004: Assessing eutrophication and reference conditions for Scottish freshwater lochs using subfossil diatoms, *Journal of Applied Ecology*, **41**: 124–138.
- Bradshaw, E.G., Rasmussen, P. and Odgaard, B.V., 2005: Mid- to late-Holocene change and lake development at Dallund Sø, Denmark: synthesis of multiproxy data, linking land and lake, *The Holocene*, **15**: 1152–1162.
- European Union, 2000: Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy, *Official Journal of the European Communities*, **L327**: 1–73.
- Haberle, S.G., Tibby, J.C., Dimitriadis, S. and Heijnis, H., 2006: The impact of European

occupation on terrestrial and aquatic ecosystem dynamics in an Australian tropical rain forest, *Journal of Ecology*, **94**: 987-1002.

**O. Heiri, M.J. Wooller, M. van Hardenbroek and Y.V. Wang**

- Brooks, S.J., 2006: Fossil midges (Diptera: Chironomidae) as palaeoclimatic indicators for the Eurasian region, *Quaternary Science Reviews*, **25**: 1894-1910.
- Cauchie, H.-M., 2002: Chitin production by arthropods in the hydrosphere, *Hydrobiologia*, **470**: 63-96.
- Grey, J., 2006: The use of stable isotope analyses in freshwater ecology: Current awareness, *Polish Journal of Ecology*, **54**: 563-584.
- Huang, Y., Shuman, B., Wang, Y. and Webb III, T., 2004: Hydrogen isotope ratios of individual lipids in lake sediments as novel tracers of climatic and environmental change: A surface sediment test, *Journal of Paleolimnology*, **31**: 363-375.
- Jones, R.I. and Grey, J., 2004: Stable isotope analysis of chironomid larvae from some Finnish forest lakes indicates dietary contribution from biogenic methane, *Boreal Environment Research*, **9**: 17-23.
- Jones, R.I., Carter, C.E., Kelly, A., Ward, S., Kelly, D.J. and Grey, J., 2008: Widespread contribution of methane-cycle bacteria to the diets of lake profundal chironomid larvae, *Ecology*, **89**: 857-864.
- Kankaala, P., Taipale, S., Grey, J., Sonninen, E., Arvola, L. and Jones, R.I., 2006: Experimental  $\delta^{13}\text{C}$  evidence for a contribution of methane to pelagic food webs in lakes, *Limnology and Oceanography*, **51**: 2821-2827.
- Lotter, A.F., Eicher, U., Birks, H.J.B. and Siegenthaler, U., 1992: Late Glacial climatic oscillations as recorded in Swiss lake sediments, *Journal of Quaternary Science*, **7**: 187-204.
- Martinez del Rio, C., Wolf, N., Carleton, S.A. and Gannes, L.Z., 2009: Isotopic ecology ten years after a call for more laboratory experiments, *Biological Reviews*, **84**: 91-111.
- Meyers, P.A. and Lallier-Vergès, E., 1999: Lacustrine sedimentary organic matter records of Late Quaternary paleoclimates, *Journal of Paleolimnology*, **21**: 345-372.
- Miller, R.F., 1991: Chitin paleoecology, *Biochemical Systematics and Ecology*, **19**: 401-411.
- Nation, J.L., 2002: *Insect physiology and biochemistry*, CRC Press, Boca Raton.
- Raabe, D., Sachs, C. and Romano, P., 2005: The crustacean exoskeleton as an example of a structurally and mechanically graded biological nanocomposite material, *Acta Materialia*, **53**: 4281-4292.
- Schimmelmann, A., accepted: Carbon, nitrogen and oxygen stable isotope ratios in chitin. In: Gupta, S.N. (Ed.), *Chitin: formation and diagenesis*, Springer.
- Schimmelmann, A., DeNiro, M.J., Poulicek, M., Voss-Foucart, M., Goffinet, G. and Jeuniaux, C., 1986: Stable isotopic composition of chitin from arthropods recovered in archaeological contexts as palaeoenvironmental indicators, *Journal of Archaeological Science*, **13**: 553-566.
- Stankiewicz, B.A., Briggs, D.E.G., Evershed, R.P. and Duncan, I.J., 1997a: Chemical preservation of insect cuticle from the Pleistocene asphalt deposit of California, USA, *Geochimica et Cosmochimica Acta*, **61**: 2247-2252.
- Stankiewicz, B.A., Briggs, D.E.G., Evershed, R.P., Flannery, M.B. and Wuttke, M., 1997b: Preservation of chitin in 25-million-year-old fossils, *Science*, **276**: 1541-1543.
- van Hardenbroek, M., Heiri, O., Grey, J., Bodelier, P.L.E., Verbruggen, F. and Lotter, A.F., 2009a: Fossil chironomid  $\delta^{13}\text{C}$  as a proxy for past methanogenic contribution to benthic food webs in lakes? *Journal of Paleolimnology*, DOI 10.1007/s10933-009-9328-5.
- van Hardenbroek, M., Heiri, O. and Lotter, A.F., 2009b: Efficiency of different mesh sizes for isolating fossil chironomids for stable isotope and radiocarbon analyses, *Journal of Paleolimnology*, DOI 10.1007/s10933-009-9327-6.

- von Grafenstein, U., Erlenkeuser, H., Brauer, A., Jouzel, J. and Johnsen, S., 1999: A mid-European decadal isotope climate record from 15,500 to 5000 years BP, *Science*, **284**: 1654-1657.
- Wang, Y., Francis, D.R., O'Brien, D.M. and Wooller, M.J., 2008: A protocol for preparing subfossil chironomid head capsules (Diptera: Chironomidae) for stable isotope analysis in paleoclimate reconstruction and considerations of contamination sources, *Journal of Paleolimnology*, **40**: 771-781.
- Wang, Y.V., O'Brien, D.M., Jenson, J., Francis, D. and Wooller, M.J., 2009: The influence of diet and water on the stable oxygen and hydrogen isotope composition of Chironomidae (Diptera) with paleoecological implications, *Oecologia*, DOI 10.1007/s00442-009-1303-3.
- Wolfe, B.B., Falcone, M.D., Clogg-Wright, K.P., Mongeon, C.L., Yi, Y., Brock, B.E., St. Amour, N.A., Mark, W.A. and Edwards, T.W.D., 2007: Progress in isotope paleohydrology using lake sediment cellulose, *Journal of Paleolimnology*, **37**: 221-231.
- Wooller, M.J., Francis, D., Fogel, M.L., Miller, U., Walker, I.R. and Wolfe, A.P., 2004: Quantitative paleotemperature estimates from  $\delta^{18}\text{O}$  of chironomid head capsules preserved in arctic lake sediments, *Journal of Paleolimnology*, **31**: 267-274.
- Wooller, M., Wang, Y. and Axford, Y., 2008: A multiple stable isotope record of Late Quaternary limnological changes and chironomid paleoecology from northeastern Iceland, *Journal of Paleolimnology*, **40**: 63-77.

#### **A. Jeziorski, A.M. Paterson, N.D. Yan and J.P. Smol**

- Alstad, N.E.W., Skardal, L. and Hessen, D.O., 1999: The effect of calcium concentration on the calcification of *Daphnia magna*, *Limnology and Oceanography*, **44**: 2011-2017.
- Ashforth, D. and Yan, N.D., 2008: The interactive effects of calcium concentration and temperature on the survival and reproduction of *Daphnia pulex* at high and low food concentrations, *Limnology and Oceanography*, **53**: 420-432.
- Cairns, A. and Yan, N., 2009: A review of the influence of low ambient calcium concentrations on freshwater daphniids, gammarids and crayfish, *Environmental Reviews*, **17**: 67-79.
- Hedin, L.O., Granat, L., Likens, G.E., Buishand, T.A., Galloway, J.N., Butler, T.J. and Rodhe, H., 1994: Steep declines in atmospheric base cations in regions of Europe and North America, *Nature*, **367**: 351-354.
- Jeffries, D.S. et al., 2003: Assessing the recovery of lakes in southeastern Canada from the effects of acidic deposition, *Ambio*, **32**: 176-182.
- Jeziorski, A. and Yan, N.D., 2006: Species identity and aqueous calcium concentrations as determinants of calcium concentrations of freshwater crustacean zooplankton, *Canadian Journal of Fisheries and Aquatic Sciences*, **63**: 1007-1013.
- Jeziorski, A. et al., 2008a: The widespread threat of calcium decline in fresh waters, *Science*, **322**: 1374-1377.
- Jeziorski, A., Paterson, A.M., Yan, N.D. and Smol, J.P., 2008b: Calcium levels in *Daphnia ephippia* cannot provide a useful paleolimnological indicator of historical lakewater Ca concentrations, *Journal of Paleolimnology*, **39**: 421-425.
- Kirchner, J.W. and Lydersen, E., 1995: Base cation depletion and potential long-term acidification of Norwegian catchments, *Environmental Science & Technology*, **29**: 1953-1960.
- Likens, G.E., Driscoll, C.T. and Buso, D.C., 1996: Long-term effects of acid rain: Response and recovery of a forest ecosystem, *Science*, **272**: 244-246.
- Peters, S.C., Lockwood, R., Williamson, C.E. and Saros, J.E., 2008: Using elemental ratios of calcium and strontium to track calcium availability in the freshwater zooplankton *Daphnia pulicaria*, *Journal of Geophysical Research*, **113**: G04023, doi:10.1029/2008JG000782.

- Smol, J.P., 2008: *Pollution of Lakes and Rivers: A Paleoenvironmental Perspective*, Second Edition, Blackwell Publishing, Malden.
- Stoddard, J.L. et al., 1999: Regional trends in aquatic recovery from acidification in North America and Europe, *Nature*, **401**: 575-578.
- Wærvågen, S.B., Rukke, N.A. and Hessen, D.O., 2002: Calcium content of crustacean zooplankton and its potential role in species distribution, *Freshwater Biology*, **47**: 1866-1878.
- Watmough, S.A., Aherne, J. and Dillon, P.J., 2003: Potential impact of forest harvesting on lake chemistry in south-central Ontario at current levels of acid deposition, *Canadian Journal of Fisheries and Aquatic Sciences*, **60**: 1095-1103.
- Watmough, S.A. and Aherne, J., 2008: Estimating calcium weathering rates and future lake calcium concentrations in the Muskoka-Haliburton region of Ontario, *Canadian Journal of Fisheries and Aquatic Sciences*, **65**: 821-833.

### **P. Rioual and L. Wang**

- An, Z.S. et al., 2006: Geophysical survey on the tectonic and sediment distribution of Qinghai Lake basin, *Science in China Series D Earth Sciences*, **49**: 851-861.
- Baier, J., Lücke, A., Negendank, J.F.W., Schleser, G.H. and Zolitschka, B., 2004: Diatom and geochemical evidence of mid- to late Holocene climatic changes at Lake Holzmaar, West-Eifel (Germany), *Quaternary International*, **113**: 81-96.
- Chen, F. et al., 2008: Holocene moisture evolution in arid central Asia and its out-of-phase relationship with Asian monsoon history, *Quaternary Science Reviews*, **27**: 351-364.
- Chu, G., Liu, J., Schettler, G., Li, J., Sun, Q., Gu, Z., Lü, H., Liu, Q. and Liu, T., 2005: Sediment fluxes and varve formation in Sihailongwan, a maar lake from northeastern China, *Journal of Paleolimnology*, **34**: 311-324.
- Chu, G., Sun, Q., Rioual, P., Boltovskoy, A., Liu, Q., Sun, P., Han, J. and Liu, J., 2008: Dinocyst microlaminations and freshwater "red tides" recorded in Lake Xiaolongwan, northeastern China, *Journal of Paleolimnology*, **39**: 319-333.
- Chu, G., Sun, Q., Gu, Z., Rioual, P., Liu, Q., Wang, K., Han, J. and Liu, J., 2009: Dust records from varved lacustrine sediments of two neighboring lakes in northeastern China over the last 1400 years, *Quaternary International*, **194**: 108-118.
- Colman, S.M., Yu, S.-Y., An, Z.S., Shen, J. and Henderson, A.C.G., 2007: Late Cenozoic climate changes in China's western interior: a review of research on Lake Qinghai and comparison with other records, *Quaternary Science Reviews*, **26**: 2281-2300.
- Cosford, J., Qing, H., Erlington, B., Matthey, D., Yuan, D., Zhang, M. and Cheng, H., 2008: East Asian monsoon variability since the Mid-Holocene recorded in a high-resolution, absolute-dated aragonite speleothem from eastern China, *Earth and Planetary Science Letters*, **275**: 296-307.
- Dearing, J.A., Jones, R.T., Shen, J., Yang, X., Boyle, J.F., Foster, G.C., Crook, D.S. and Elvin, M.J.D., 2008: Using multiple archives to understand past and present climate-human-environment interactions: the lake Erhai catchment, Yunnan Province, China, *Journal of Paleolimnology*, **40**: 3-41.
- Dong, X., Bennion, H., Battarbee, R.W., Yang, X., Yang, H. and Liu, E. 2008: Tracking eutrophication in Taihu Lake using diatom record: potential and problems, *Journal of Paleolimnology*, **40**: 413-429.
- Ji, J., Balsam, W., Shen, J., Wang, M., Wang, H. and Chen, J., 2009: Centennial blooming of anoxygenic phototrophic bacteria in Qinghai Lake linked to solar and monsoon activities during the last 18,000 years, *Quaternary Science Reviews*, doi.10.1016/j.quascirev.2008.12.015
- Liu, J., Liu, T. and Negendank, J.F.W., 2001: The Chinese maar drilling programme. A Chinese-German cooperation for paleoclimatic reconstructions, *Pages News*, **9**: 10-11.

- Liu, Z., Henderson, A.C.G. and Huang, Y., 2006: Alkenone-based reconstruction of late-Holocene surface temperature and salinity changes in Lake Qinghai, China, *Geophysical Research Letters*, **33**: L09707, doi:10.1029/2006GL026151.
- Liu, J., Chu, G., Han, J., Rioual, P., Jiao, W. and Wang, K., 2009: Volcanic eruptions in the Longgang volcanic field, northeastern China, during the past 15,000 years, *Journal of Asian Earth Sciences*, **34**: 645-654.
- Mingram, J. et al., 2007: East Asian monsoon variability over the last 65,000 years – new insights from laminated lake sediments of NE-China. 3rd Alexander von Humboldt International conference, *The East Asian Monsoon, past, present and future*, Beijing, China, Abstract volume p14.
- Parplies, J., Lücke, A., Vos, H., Mingram, J., Stebich, M., Radtke, U., Han, J. and Schleser, G.H., 2008: Late glacial environment and climate development in northeastern China derived from geochemical and isotopic investigations of the varved sediment record from Lake Sihailongwan (Jilin Province), *Journal of Paleolimnology*, **40**: 471-487.
- Schettler, G., Mingram, J., Negendank, J.F.W. and Liu, J., 2006a: Palaeovariations in the East-Asian monsoon regime geochemically recorded in varved sediments of Lake Sihailongwan (Northeast China, Jilin Province). Part 2: a 200-year record of atmospheric lead-210 flux variations and its palaeoclimatic implications, *Journal of Paleolimnology*, **35**: 271-288.
- Schettler, G., Liu, Q., Mingram, J., Stebich, M. and Dulski, P., 2006b: East-Asian monsoon variability between 15 000 and 2000 cal. yr BP recorded in varved sediments of Lake Sihailongwan (northeastern China, Long Gang volcanic field), *The Holocene*, **16**: 1043-1057.
- Stebich, M., Mingram, J., Han, J. and Liu, J., 2009: Late Pleistocene spread of (cool-) temperate forests in Northeast China and climate changes synchronous with the North Atlantic region, *Global and Planetary Change*, **65**: 56-70.
- Shen, J., Jones, R.T., Yang, X., Dearing, J.A. and Wang, S., 2006: The Holocene vegetation history of Lake Erhai, Yunnan province southwestern China: the role of climate and human forcings, *The Holocene*, **16**: 265-276.
- Wang, L. et al., 2008: Diatom-based inference of variations in the strength of Asian winter monsoon winds between 17,500 and 6000 calendar years B.P., *Journal of Geophysical Research*, **113**: D21101, doi:10.1029/2008JD010145.
- Wang, S., Lü, H., Liu, J. and Negendank, J.F.W., 2007: The early Holocene optimum inferred from a high-resolution pollen record of Huguanyan Maar Lake in southern China, *Chinese Science Bulletin*, **52**: 2829-2836.
- Wang, Y. et al., 2005: The Holocene Asian monsoon: links to solar changes and North Atlantic climate, *Science*, **308**: 854-857.
- Xiao, J., Xu, Q., Nakamura, T., Yang, X., Liang, W. and Inouchi, Y., 2004: Holocene vegetation variation in the Dahai Lake region of north-central China: a direct indication of the Asian monsoon climatic history, *Quaternary Science Reviews*, **23**: 1669-1679.
- Xiao, J., Si, B., Zhai, D., Itoh, S. and Lomtatidze, Z., 2008: Hydrology of Dali Lake in central-eastern Inner Mongolia and Holocene East Asian monsoon variability, *Journal of Paleolimnology*, **40**: 519-528.
- Yancheva, G. et al., 2007: Influence of the intertropical convergence zone on the East Asian monsoon, *Nature*, **445**: 74-77.
- Yang, X., Anderson, N.J., Dong, X. and Shen, J., 2008: Surface sediment diatom assemblages and epilimnetic total phosphorus in large, shallow lakes of the Yangtze floodplain: their relationships and implications for assessing long-term eutrophication, *Freshwater Biology*, **53**: 1273-1290.
- You, H., Liu, J., Liu, Q., Chu, G., Rioual, P. and Han, J., 2008: Study of the varve record from Erlongwan maar lake, NE China, over the last 13 ka BP, *Chinese Science Bulletin*, **53**: 262-266.

Zhang, D. and Lu, L., 2007: Anti-correlation of summer/winter monsoons? *Nature*, **450**: E7-E9.

**P. Rosén, H. Vogel, L. Cunningham, N. Reuss, D. Conley and P. Persson**

Calace, N., Capolei, M., Lucchese, M. and Petronio, B.M., 1999: The structural composition of humic compounds as indicator of organic carbon sources, *Talanta*, **49**: 277-284.

Mecozzi, M. and Pietrantonio, E., 2006: Carbohydrates proteins and lipids in fulvic and humic acids of sediments and its relationships with mucilaginous aggregates in the Italian seas, *Marine Chemistry*, **101**: 27-39.

Müller, P.J. and Schneider, J., 1993: An automated leaching method for the determination of opal in sediments and particulate matter, *Deep-Sea Research*, **40**: 425-444.

Rosén, P. and Persson, P., 2006: Fourier-transform infrared spectroscopy (FTIRS), a new method to infer past changes in tree-line position and TOC using lake sediment, *Journal of Paleolimnology*, **35**: 913-923.

Rosén, P., Vogel, H., Cunningham, L., Reuss, N., Conley, D. and Persson, P., 2009: Fourier transform infrared spectroscopy, a new method for rapid determination of total organic and inorganic carbon and opal concentration in lake sediments, *Journal of Paleolimnology*, DOI: 10.1007/s10933-009-9329-4.

Schnürer, Y., Persson, P., Nilsson, M., Nordgren, A. and Giesler, R., 2006: Effects of surface sorption on microbial degradation of glyphosate, *Environmental Science and technology*, **40**: 4145-4150.

Stehfest, K., Toepel, J. and Wilhelm, C., 2005: The application of micro-FTIR spectroscopy to analyze nutrient stress-related changes in biomass composition of phytoplankton algae, *Plant Physiology and Biochemistry*, **43**: 717-726.

Vogel, H., Rosén, P., Wagner, B., Melles, M. and Persson, P., 2008: Fourier transform infrared spectroscopy, a new cost-effective tool for quantitative analysis of biogeochemical properties in long sediment records, *Journal of Paleolimnology*, **40**: 689-702.

White, W.B., 1974: The carbonate minerals, p. 227-242. In: Farmer, V.C. (ed.), 1974. The infrared spectra of minerals. *Mineralogical Society Monograph* 4.

**P. Tarasov, S. Müller, A. Andreev, K. Werner and B. Diekmann**

Abbott, R.J., Smith, L.C., Milne, R.I., Crawford, R.M.M., Wolff, K. and Balfour, J., 2000: Molecular analysis of plant migration and refugia in the Arctic, *Science*, **289**: 1343-1346.

Binney, H.A., Willis, K.J., Edwards, M.E., Bhagwat, S.A., Anderson, P.M., Andreev, A.A., Blaauw, M., Damblon, F., Haesaerts, P., Kienast, F., Kremenetski, K.V., Krivonogov, S.K., Lozhkin, A.V., MacDonald, G.M., Novenko, E.Y., Oksanen, P., Sapelko, N.V., Väliranta, M. and Vazhenina, L., 2009: The distribution of late-Quaternary woody taxa in northern Eurasia: evidence from a new macrofossil database, *Quaternary Science Reviews*, **28**: 2445-2464.

Brubaker, L.B., Anderson, P.M., Edwards, M.E. and Lozhkin, A.V., 2005: Beringia as a glacial refugium for boreal trees and shrubs: new perspectives from mapped pollen data, *Journal of Biogeography*, **32**: 833-848.

Danzeglocke, U., Jöris, O. and Weninger, B., 2008: CalPal-2007online. <http://www.calpal-online.de>.

Johnsen, S.J., Dahl-Jensen, D., Gundestrup, N., Steffensen, J.P., Clausen, H.B., Miller, H., Masson-Delmotte, V., Sveinbjörnsdottir, A.E. and White, J., 2001: Oxygen isotope and palaeotemperature records from six Greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP, *Journal of Quaternary Science*, **16**: 299-307.

MacDonald G.M., 2001: Conifer stomata. In: Smol J.P., Birks H.J.B. and Last W.M. (eds), *Tracking Environmental Change Using Lake Sediments. Vol. 3: Terrestrial, Algal and*

Siliceous Indicators. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 33–47.

- MacDonald, G.M., Velichko, A.A., Kremenetski, C.V., Borisova, O.K., Goleva, A.A., Andreev, A.A., Cwynar, L.C., Riding, R.T., Forman, S.L., Edwards, T.W.D., Aravena, R., Hammarlund, D., Szeicz, J.M. and Gattaulin, V.N., 2000: Holocene treeline history and climate change across Northern Eurasia, *Quaternary Research*, **53**: 302–311.
- Müller, S., Tarasov, P.E., Diekmann, B. and Andreev, A.A., 2009: Late Glacial to Holocene environments in the present-day coldest region of the Northern Hemisphere inferred from a pollen record of Lake Billyakh, Verkhoyansk Mts, NE Siberia, *Climate of the Past*, **5**: 73-84.
- Petit, R.J., Hu, F.S. and Dick, C.W., 2008: Forests of the Past: A Window to Future Changes, *Science*, **320**: 1450-1452.
- Prentice, I.C., Crameer, W., Harrison, S.P., Leemans, R., Monserud, R.A. and Solomon, A.M., 1992: A global biome model based on plant physiology and dominance, soil properties, and climate, *Journal of Biogeography*, **19**: 117–134.
- Tarasov, P., Williams, J.W., Andreev, A., Nakagawa, T., Bezrukova, E., Herzs Schuh, U., Igarashi, Y., Müller, S., Werner, K. and Zheng, Z., 2007: Satellite- and pollen-based quantitative woody cover reconstructions for northern Asia: verification and application to late-Quaternary pollen data, *Earth and Planetary Science Letters*, **264**: 284–298.
- Werner, K., Tarasov, P.E., Andreev, A.A., Müller, S., Kienast, F., Zech, M., Zech, W. and Diekmann, B., 2009: A 12.5-kyr history of vegetation dynamics and mire development with evidence of Younger Dryas larch presence in the Verkhoyansk Mountains, East Siberia, Russia, *Boreas*, published online.

#### **D. Verschuren and J.M. Russell**

- Bergner, A.G.N., Trauth, M.H. and Bookbagen, B., 2003: Paleoprecipitation estimates for the Lake Naivasha basin (Kenya) during the last 175 ky using a lake-balance model, *Global and Planetary Change*, **36**: 117-136.
- Castañeda, I.S., Werne, J.P. and Johnson, T.C., 2007: Wet/arid phases in the southeast African tropics since the Last Glacial Maximum, *Geology*, **35**, 823-826.
- Cohen, A.S., Palacios-Fest, M.R., Msaky, E.S., Alin, S.R., McKee, B.A., O'Reilly, C.M., Dettman, D.L., Nkotagu, H., and Lezzar, K.E., 2005: Paleolimnological investigations of anthropogenic environmental change in Lake Tanganyika: IX. Summary of paleorecords of environmental change and catchment deforestation at Lake Tanganyika and impacts on the Lake Tanganyika ecosystem, *Journal of Paleolimnology*, **34**: 125-145.
- Cohen, A.S., et al., 2007: Ecological consequences of early Late Pleistocene mega-droughts in tropical Africa, *Proceedings of the National Academy of Science USA*, **104**: 16422-16427.
- Duhnforth, M., Bergner, A.G.N. and Trauth, M.H., 2006: Early holocene water budget of the Nakuru-Elmenteita Basin, Central Kenya Rift, *Journal of Paleolimnology*, **36**: 281-294.
- Eggermont, H., Heiri, O. M., Russell, J. M., Vuille, M., Audenaert, L. and Verschuren, D., 2009: Paleotemperature reconstruction in tropical Africa using fossil Chironomidae (Insecta: Diptera), *Journal of Paleolimnology*, doi 10.1007/s10933-009-9339-2.
- Eggermont, H., Heiri, O. M. and Verschuren, D., 2006: Subfossil Chironomidae (Insecta: Diptera) as quantitative indicators of past salinity variation in African lakes, *Quaternary Science Reviews*, **25**: 1966-1994.
- Felton, A.A., et al., 2007: Paleolimnological evidence for the onset and termination of glacial aridity from Lake Tanganyika, Tropical East Africa, *Palaeogeography, Palaeoclimatology, Palaeoecology*, **252**: 405-423.

- Fleitmann, D., Dunbar, R.B., McCulloch, M., Mudelsee, M., Vuille, M., McClanahan, T., Andrews, C. and Mucciarone, D.A., 2007: East African soil erosion recorded in a 300-year old coral colony from Kenya, *Geophysical Research Letters*, **34**: L04401.
- Garcin, Y., Vincens, A., Williamson, D., Guiot, J. and Buchet, G., 2006: Wet phases in tropical southern Africa during the last glacial period, *Geophysical Research Letters*, **33**: L07703.
- Garcin, Y., Vincens, A., Williamson, D., Buchet, G. and Guiot, J., 2007: Abrupt resumption of the African Monsoon at the Younger Dryas-Holocene climatic transition, *Quaternary Science Reviews*, **26**: 690-704.
- Gasse, F., Chalié, F., Vincens, A., Williams, M.A.J. and Williamson, D., 2008: Climatic patterns in equatorial and southern Africa from 30,000 to 10,000 years ago reconstructed from terrestrial and near-shore proxy data, *Quaternary Science Reviews*, **27**: 2316-2340.
- Johnson, T.C., Brown, E.T., McManus, J., Barry, S., Barker, P. and Gasse, F., 2002: A high-resolution paleoclimate record spanning the past 25,000 years in southern East Africa, *Science*, **296**: 113-116.
- Kawamura, K. et al., 2007: Northern hemisphere forcing of climatic cycles in Antarctica over the past 360,000 years, *Nature*, **448**: 912-916.
- Kröpelin, S. et al., 2008: Climate-driven ecosystem succession in the Sahara: the past 6000 years, *Science*, **320**: 765-768.
- Lamb, H., Darbyshire, L. and Verschuren, D., 2003: Vegetation response to rainfall variation and human impact in central Kenya during the past 1100 years, *The Holocene*, **13**: 285-292.
- Lamb, H.F., Leng, M.J., Telford, R.J., Ayenew, T. and Umer, M., 2007. Oxygen and carbon isotope, composition of authigenic carbonate from an Ethiopian lake: a climate record of the last 2000 years, *The Holocene*, **17**, 517-526.
- Legesse, D., Vallet-Coulomb, C. and Gasse, F., 2004: Analysis of the hydrological response of a tropical terminal lake, Lake Abiyata (Main Ethiopian Rift Valley) to changes in climate and human activities, *Hydrological Processes*, **18**, 487-504.
- Lejju, J.B., 2009: Vegetation dynamics in western Uganda during the last 1000 years: climate change or human induced environmental degradation? *African Journal of Ecology*, **47**: 21-29.
- Mergeay, J., Verschuren, D., Van Kerckhoven, L. and De Meester, L., 2004: Two hundred years of a diverse *Daphnia* community in Lake Naivasha (Kenya): effects of natural and human-induced environmental changes, *Freshwater Biology*, **49**: 998-1013.
- Mergeay, J., Verschuren, D. and De Meester, L., 2006: Invasion of an asexual American water flea clone throughout Africa and rapid displacement of a native sibling species, *Proceedings of the Royal Society London B*, **273**: 2839-2844.
- Mergeay, J., Vanoverbeke, J., Verschuren, D. and De Meester, L., 2007: Extinction, re-colonization, and dispersal through time in a planktonic crustacean, *Ecology*, **88**: 3032-3043.
- Nash, D.J., Meadows, M.E. and Gulliver, V.L., 2006: Holocene environmental change in the Okavango Panhandle, northwest Botswana, *Quaternary Science Reviews*, **25**: 1302-1322.
- Ngomanda, A., Neumann, K., Schweizer, A. and Maley, J., 2009: Seasonality change and the third millennium BP rainforest crisis in southern Cameroon (Central Africa), *Quaternary Research*, **71**: 307-318.
- O'Reilly, C.M., Alin, S.R., Plisnier, P.D., Cohen, A.S. and McKee, B.A., 2003: Climate change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa, *Nature*, **424**: 766-768.
- Powers, L. A., Johnson, T.C., Werne, J.P., Castañeda, I.S., Hopmans, E.F., Sinninghe Damsté, J.S. and Schouten, S., 2005: Large temperature variability in the southern African tropics since the Last Glacial Maximum, *Geophysical Research Letters*, **32**: L08706.

- Rumes, B., Eggermont, H. and Verschuren, D., 2005: Subfossil representation of aquatic invertebrate communities in a salinity gradient of western Uganda crater lakes, *Hydrobiologia*, **542**: 297-314.
- Russell, J.M. and Johnson, T.C., 2007. Little Ice Age drought in equatorial Africa: ITCZ migrations and ENSO variability, *Geology*, **35**: 21-24.
- Russell, J.M., McCoy S.J., Verschuren, D., Bessems, I., Blaauw, M. and Huang, Y., 2009: Human impacts, climate change, and aquatic ecosystem response during the past 2,000 years at Lake Wandakara, Uganda, *Quaternary Research*, doi 10.1016/j.yqres.2009.06.008.
- Scholz, C.A., et al., 2007: East African mega-droughts between 135 and 75 thousand years ago and bearing on early-modern human origins, *Proceedings of the National Academy of Sciences USA*, **104**: 16416-16421.
- Ssemmanda, I., Ryves, D.B., Bennike, O. and Appleby, P.G., 2005: Vegetation history in western Uganda during the last 1200 years: a sediment-based reconstruction from two crater lakes, *The Holocene*, **15**: 119-132.
- Shanahan, T.M., Overpeck, J.T., Anchukaitis, K.J., Beck, J.W., Cole, J.E., Dettman, D.L., Peck, J.A., Scholz, C.A. and King, J.W., 2009: Atlantic forcing of persistent drought in West Africa, *Science*, **324**: 377-380.
- Stager, J.C., Ryves, D., Cumming, B.F., Meeker, L.D. and Beer, J., 2005: Solar variability and the levels of Lake Victoria, East Africa, during the last millennium, *Journal of Paleolimnology*, **33**: 243-251.
- Talbot, M.R., Filippi, M.L., Jensen, N.B. and Tiercelin, J.-J., 2007: An abrupt change in the African monsoon at the end of the Younger Dryas, *Geochemistry, Geophysics, Geosystems*, **8**: Q03005.
- Tierney, J.E., Russell, J.M., Huang, Y.S. and Cohen, A.S., 2008: Northern Hemisphere controls on tropical Southeast African climate during the last 60,000 years, *Science*, **322**: 252-255.
- Verschuren, D. and Charman, D., 2008: Latitudinal linkages in late-Holocene moisture-balance variation. In: Battarbee, R.W. et al. (Eds), *Natural climate variability and global warming*, Blackwell, 189-231.
- Verschuren, D., Tibby, J., Leavitt, P.R. and Roberts, C.N., 1999: The environmental history of a climate-sensitive lake in the former 'White Highlands' of central Kenya, *Ambio*, **28**: 494-501.
- Verschuren, D., Laird, K.R. and Cumming, B.F., 2000a: Rainfall and drought in equatorial East Africa during the past 1100 years, *Nature*, **403**: 410-414.
- Verschuren, D., Tibby, J., Sabbe, K. and Roberts, N., 2000b: Effects of lake level, salinity and substrate on the invertebrate community of a fluctuating tropical lake, *Ecology* **81**: 164-182.
- Verschuren, D., Johnson, T.C., Kling, H.J., Edgington, D.N., Leavitt, P.R., Brown, E.T., Talbot, M.R. and Hecky, R.E., 2002: History and timing of human impact on Lake Victoria, East Africa, *Proceedings of the Royal Society London B*, **269**: 289-294.

**J.W.H. Weijers, C.I. Blaga, J.P. Werne and J.S. Sinninghe Damsté**

- Blaga, C.I., Reichart, G.-J., Heiri, O. and Sinninghe Damsté, J.S. 2009: Tetraether membrane lipid distributions in water-column particulate matter and sediments: a study of 47 European lakes along a north-south transect, *Journal of Paleolimnology*, **41**: 523-540.
- Kim J.-H., Schouten S., Hopmans E.C., Donner B. and Sinninghe Damsté J.S., 2008: Global sediment core-top calibration of the TEX<sub>86</sub> paleothermometer in the ocean, *Geochimica et Cosmochimica Acta*, **72**: 1154-1173.
- Lotter, A.F., Birks, H.J.B., Hofmann, W. and Marchetto, A., 1997: Modern diatom, cladocera, chironomid and chrysophyte cyst assemblages as quantitative indicators for the

- reconstruction of past environmental conditions in the Alps. I. Climate, *Journal of Paleolimnology*, **18**: 395-420.
- Peterse, F., Kim, J.-H., Schouten, S., Klitgaard-Kristensen, D., Koç, N. and Sinninghe Damsté, J.S., 2009: Constraints on the application of the MBT/CBT paleothermometer at high latitude environments (Svalbard, Norway), *Organic Geochemistry*, **40**: 692-699.
- Powers, L.A., Werne, J.P., Johnson, T.C., Hopmans, E.C., Sinninghe Damsté, J.S. and Schouten, S., 2004: Crenarchaeotal membrane lipids in lake sediments; a new paleotemperature proxy for continental paleoclimate reconstruction? *Geology*, **32**: 613-616.
- Powers, L.A., Johnson, T.C., Werne, J.P., Castañeda, I.S., Hopmans, E.C., Sinninghe Damsté, J.S. and Schouten, S., 2005: Large temperature variability in the southern African tropics since the Last Glacial Maximum, *Geophysical Research Letters*, **32**: L08706, doi:10.1029/2004GL022014.
- Riatti-Shanti, M., Shemesh, A. and Karlen, W., 1998: A 3000-year climatic record from biogenic silica oxygen isotopes in an equatorial high altitude lake, *Science*, **281**: 980-982.
- Schouten, S., Hopmans, E.C., Schefuss, E. and Sinninghe Damsté, J.S., 2002: Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water temperatures? *Earth and Planetary Science Letters*, **204**: 265-274.
- Sinninghe Damsté, J.S., Ossebaar, J., Abbas, B., Schouten, S. and Verschuren, D., 2009: Fluxes and distribution of tetraether lipids in an equatorial African lake: Constraints on the application of the TEX<sub>86</sub> palaeothermometer and BIT index in lacustrine settings, *Geochimica et Cosmochimica Acta*, **73**: 4232-4249.
- Sluijs, A. and Schouten, S. et al., 2006: Subtropical Arctic Ocean temperatures during the Palaeocene/Eocene thermal maximum, *Nature*, **441**: 610-613.
- Tierney, J.E., Russel, J.M., Huang, Y., Sinninghe Damsté, J.S., Hopmans, E.C. and Cohen, A.S., 2008: Northern Hemisphere controls on tropical Southeast African climate during the past 60,000 years, *Science*, **322**: 252-255.
- Tierney, J.E. and Russel, J.M., 2009: Distributions of branched GDGTs in a tropical lake system: implications for lacustrine application of the MBT/CBT paleoproxy, *Organic Geochemistry*, **40**: 1032-1036.
- von Grafenstein, U., Erlenkeuser, H., Kleinmann, J., Müller, J. and Trumborn, P., 1994: High frequency climatic oscillations during the last deglaciation as revealed by oxygen-isotope records of benthic organisms (Ammersee, southern Germany), *Journal of Paleolimnology*, **11**: 349-357.
- Wakeham, S.G., Lewis, C.M., Hopmans, E.C., Schouten, S. and Sinninghe Damsté, J.S., 2003: Archaea mediate anaerobic oxidation of methane in deep euxinic waters of the Black Sea, *Geochimica et Cosmochimica Acta*, **67**: 1359-1374.
- Walker, I.R., Smol, J.P., Engstrom, D.R. and Birks, H.J.B., 1991: An assessment of Chironomidae as quantitative indicators of past climatic change, *Canadian Journal of Fisheries and Aquatic Science*, **48**: 975-987.
- Weijers, J.W.H., Schouten, S., Spaargaren, O.C. and Sinninghe Damsté, J.S., 2006: Occurrence and distribution of tetraether membrane lipids in soils: Implications of the use of the TEX<sub>86</sub> proxy and the BIT index, *Organic Geochemistry*, **37**: 1680-1693.
- Weijers, J.W.H., Schouten, S., van den Donker, J.C., Hopmans, E.C. and Sinninghe Damsté, J.S., 2007a: Environmental controls on bacterial tetraether membrane lipid distribution in soils, *Geochimica et Cosmochimica Acta*, **71**: 703-713.
- Weijers, J.W.H., Schefuß, E., Schouten, S., Sinninghe Damsté, J.S., 2007b: Coupled thermal and hydrological evolution of tropical Africa over the last deglaciation, *Science*, **315**: 1701-1704.

Weijers, J.W.H., Schouten, S., Sluijs, A., Brinkhuis, H. and Sinninghe Damsté, J.S., 2007c:  
Warm arctic continents during the Palaeocene-Eocene thermal maximum, *Earth and  
Planetary Science Letters*, **261**: 230-238.